

PUPPY SHOT CALENDAR

Puppy / Litter Name : _____

Date Bred :

Open Eyes :

Whelp Date :

Can Hear :

Wormer

Wormer should be started when puppies are two weeks old, then given every 2 weeks after, until 12 weeks old, then switched to once a month until 6 months old. At 6 months old, dogs can be put on adult wormer.

- | | | | |
|---|---|---|--|
| <input type="checkbox"/> Pyrantel | <input type="checkbox"/> Fenbendazole | <input type="checkbox"/> Milbemycin, Moxidectin, Ivermectin (Adult) | |
| <input type="checkbox"/> 2 Weeks _____ | <input type="checkbox"/> 4 Weeks _____ | <input type="checkbox"/> 6 Weeks _____ | <input type="checkbox"/> 8 Weeks _____ |
| <input type="checkbox"/> 10 Weeks _____ | <input type="checkbox"/> 12 Weeks _____ | <input type="checkbox"/> +1 months _____ | <input type="checkbox"/> +2 months _____ |

Vaccination

These base vaccine protect against Distemper, Hepatitis/Adenovirus, Parainfluenza, Parvovirus. A minimum of 3 doses, with first dose starting at 6 weeks old, then given every 2-4 weeks after. Can be administered in the loose skin between the shoulder blades.

- | | | | |
|--|--|--|---|
| <input type="checkbox"/> 4-in-1 DHPP | <input type="checkbox"/> 5-in-1 - DHHPP | <input type="checkbox"/> 6-in-1 - DHHCPP | <input type="checkbox"/> 7-in-1 - DHHPLL |
| <input type="checkbox"/> 6-8 Weeks _____ | <input type="checkbox"/> 10-12 Weeks _____ | <input type="checkbox"/> 14-16 Weeks _____ | <input type="checkbox"/> Annual Booster _____ |

Bordetella

This shot protects against kennel cough, a contagious respiratory infection. Should be started at 6-8 weeks old, then given again in 2-4 weeks. Can be administered in the loose skin between the shoulder blades.

- | | |
|--|--|
| <input type="checkbox"/> 6-8 Weeks _____ | <input type="checkbox"/> 10-12 Weeks _____ |
|--|--|

Canine Lyme Disease

This shot protects against Lyme Disease, a tick-borne illness that can lead to fever, joint pain, kidney and heart problems. Should be started at 10-12 weeks old, then given again in 2-4 weeks

- | | |
|--|--|
| <input type="checkbox"/> 10-12 Weeks _____ | <input type="checkbox"/> 14-16 Weeks _____ |
|--|--|

Canine Influenza

This vaccine protects your dog against canine influenza, a contagious respiratory infection that can progress into pneumonia. Should be started at 10-12 weeks old, then given again in 2-4 weeks

- | | |
|--|--|
| <input type="checkbox"/> 10-12 Weeks _____ | <input type="checkbox"/> 14-16 Weeks _____ |
|--|--|

Leptospirosis

This shot protects against, a bacterial disease spread through infected urine. Maybe included in some 5-in-1 shots. Should be given at 12 weeks old, then again in 2-4 weeks. Can be given in the loose skin between the shoulder blades or side of neck behind the shoulders.

- | | |
|---|--|
| <input type="checkbox"/> 12 Weeks _____ | <input type="checkbox"/> 14-16 Weeks _____ |
|---|--|

Rabies

This shot protects against rabies, an uncommon, but deadly viral disease that effects all mammals. Should be given at 14-16 weeks old and may be required by law.

- | | |
|---|------------|
| <input type="checkbox"/> 14 Weeks _____ | ID # _____ |
|---|------------|

Rattlesnake toxoid

This shot protects against rattlesnake venom. Maybe recommended by your vet if you live in areas with heavy rattlesnake activity.

- | |
|---|
| <input type="checkbox"/> 14 Weeks _____ |
|---|

